

The Cytoprotective and Cell Recovery Properties of Apple Extracts on H₂O₂ induced-NIH3T3 Cells : An Anti Aging Candidate

Nunuk Aries Nurulita^{1,2,*}, Anjar Mahardian Kusuma¹, Darsini¹, Weny Delvia¹, Veby Tri Yulianti¹

¹Faculty of Pharmacy, Universitas Muhammadiyah Purwokerto, Purwokerto, Indonesia

²Cancer and Stem Cell Research Center (CSCR), Universitas Muhammadiyah Purwokerto, Purwokerto, Indonesia

Abstract

Apple contains high concentration of phenolic compounds that protect cells from oxidative stress. The prolong exposure of free radicals may induce cell damage and premature cell aging. Both local and imported apple contain flavonoid, saponin, tannin, steroid, and terpenoid. The extract of local and imported apples showed low toxicity on NIH3T3 fibroblast cells, with IC₅₀ value of 529 and 463 µg/mL, respectively. Both apple extracts (50-250 µg/mL) protected three-day-H₂O₂ induced-cell damage and cell death. Protective effect was observed as the viability increase of treated cells compared to untreated ones. The protective effect of both extracts were higher than the effect of vitamin C as standard antioxidant at this study. Both apple extracts could reverse cell damage caused by three-hour-high concentration H₂O₂ exposure, similar with vitamin C. Low concentration of both extracts (50 µg/mL) induced the increase of fibroblast cells' proliferation kinetics. The extract of imported apple showed higher properties of protective, cell recovery and proliferation of fibroblast cells than local apple, but not statistically significance. This study concludes that the extract of local and imported apples have high potency in cytoprotective effect and cell recovery of damaged cells caused by free radicals induction. Both apple extracts have high potency to be developed the candidate of antiaging and cells' regeneration agent.

Keywords : *Antiaging, cell recovery, cytoprotective, NIH3T3 cells*

INTRODUCTION

Aging is one natural process occurring during living organism's life. It is a complex process causing progressive decrease in physiological function, followed by dysfunction, and death. Several factors play a role in growth and aging, that are disease, injury, nutrition intake, exercises, stress, and plenty of environmental factors (Curtis, *et al.*, 2005). Skin roughness and dryness followed

by loss of flexibility and fairness are also the signs of skin aging (Atmaja, *et al.*, 2012). Scientific efforts have been done to slow aging process by lengthening healthy age duration (Getoff, 2007).

Submitted: June 6, 2018

Revised: June 10, 2018

Accepted: June 30, 2018

*Corresponding author: nunukariesnurulita@ump.ac.id

Corrective treatment or repair could be done by using microdermabrasion, iontophoresis, laser, chemical peeling, mesotherapy, sonophoresis, and application of night cream or skin peeling cream that helps reducing fleck. It drives people's interest in improving ways to slow aging process (Gilchrest, *et al.*, 2003).

Kujawa, *et al.*, (2011) observed that apple peel extract possessed high antioxidant activity, almost comparable to that of its fruit's extract. Pertiwi, *et al.* (2016) also proved antioxidant activity of apple peel's ethanolic extract, giving IC₅₀ of 87 ppm. Juranovic, *et al.*, (2011) showed that apple extract has antiaging activity by observing yeast cells viability *in vitro*. Yuniarto, *et al.* (2012) revealed that apple fruit's extract possessed antioxidant activity with IC₅₀ of 151 ppm. Medium-sized fresh apple extract contains antioxidants similar to that of 1,500 mg of vitamin C (Nurcahyati, 2014). Quercetin, known as the flavonol mostly found in nature compared to other flavonoids, is contained in apple fruit, believed to be responsible for its antioxidant and antiaging activity (Wasim and Farhan, 2010).

A study on the antioxidant activity of a set of Indonesian's fruits showed that flavonoids contained in peel extracts of rambutan, durian, sweet orange, kelengkeng, and kelengkeng seed were 12.26; 46.03; 9.34; 8.82; and 5.21 mg/g, respectively. In this study, we observed cytoprotective activity of both local and imported apple extract against H₂O₂ exposure on NIH3T3 fibroblast cells, since fibroblast cells are human's cell commonly used in wound healing and early aging studies.

MATERIALS AND METHODS

Materials

Materials used in this study were local apples (Malang green apple, Indonesia), imported apples (Thailand green apple), ethanol 70% (Brataco), NIHT3T fibroblast cell line, DMEM powder, Fetal Bovine Serum (FBS) (Sigma), penicillin-streptomycin (Sigma), ascorbic acid, 3% H₂O₂

(Kimia Farma), PBS, MTT (Calbiochem), Trypsin (Sigma), and Dimethyl Sulfoxide (DMSO) (Sigma). Centrifuge, CO₂ incubator (Thermo Scientific), inverted microscope (Olympus), conicals (Biologics), ELISA reader (Bio-rad iMark™), and LAF (Thermo Scientific) were used.

Apple Extraction

Apple dried simplicia was extracted using 70% ethanol by three-cycle maceration for 5 days in each cycle. Macerate obtained was concentrated using rotary evaporator to give concentrated extract.

Screening of Secondary Metabolites

Identification was conducted by specific color-forming reactions suitable for each group of substance. 2 NHCl was used for flavonoid (glycoside-3-flavonols) identification, Meyer reagent was used for alkaloids identification, FeCl₃ was used for tannins identification, foam-formation reaction was used for saponins identification, and glacial acetic acid together with concentrated sulfuric acid were used for terpenoids and steroids identification.

Cell Viability Assay (MTT Assay)

As much as 5 x 10³ cell/well of NIH3T3 cells were grown in 96-well plate and incubated with 5% CO₂ flow in 37°C. Media was removed and cells were washed with 100 µL PBS. For cytotoxicity assay, 0-1000 ppm of apple extract were applied for 24 hours, while for proliferative assay 10, 25, 50, and 125 ppm of apple extract were applied for 0, 24, 48, and 72 hours in normal growth condition (37°C, 5% CO₂ flow). Following treatments, culture media was removed. PBS washing was done prior to 100 µL MTT reagent addition followed by 4 hours incubation (37°C, 5% CO₂ flow) to ensure complete formazan crystals formation). As much as 100 µL SDS in HCl was added to each well, and cells were incubated overnight, followed by absorbance measurement using ELISA reader at 595 nm wavelength.

Induction and Stress Recovery Testing using H₂O₂

As much as 5 x 10³ cells/well were grown in 96 well plate, and were incubated for 24 hours in 37°C with 5% CO₂ flow. Culture media was removed, and cells were washed with 100 µL PBS. Treatment of 0-1000 ppm apple extract and 325 ppm ascorbic acid as positive control were given. After 1 hour, 100 µL H₂O₂ was added to each well for 1 hour, followed by media replacement with fresh media. The treatment were given in 3 consecutive days to induce stress and cell recovery following H₂O₂ exposure. After that, MTT was added to each well, and cells were incubated for 4 hours (37°C, 5% CO₂ flow). As much as 100 µL SDS was added prior to overnight incubation, and absorbance measurement using ELISA reader at 595nm wavelength.

Data Analysis

Cytotoxicity assay was analyzed by IC₅₀ value calculation using Probit analysis (SPSS 16 for Windows software). The differences among groups were analyzed using Post-hoc Tukey's two-way Anova and Kruskal Wallis followed by Mann-Whitney test.

RESULTS AND DISCUSSION

Apple Fruit Extraction

Extraction of local and imported apple through maceration gave quite high yield (Table 1). Since ethanol is a universal solvent that is semipolar, most

Table 1. LAE and IAE extracts.

Sample	Yield (%)	Organoleptic	
		Color	Smell
Local Apple	11.835	Dark brown	Typical unique
Imported Apple	10.587	Dark brown	Typical unique

of the substances contained in apple was expected to be extracted. The difference of yield obtained from the two extracts could possibly be caused by the difference of secondary metabolites content between them. It came as the result of the difference of growth territory, weather, and nutrition content in the soil where the plant grew.

Secondary Metabolites Screening

Secondary metabolites screening results showed that both local and imported apples contained flavonoids, saponins, tannins, alkaloids, steroids, and terpenoids (Table 2).

Apple Extracts Cell Viability Testing by MTT Assay

Local Apple extract (LAE) and Imported Apple extract (IAE) treatment (0-1000 µg/mL) brought change in NIH3T3 fibroblast cells morphology. Fibroblast cells became rounder, darken, and some cells were observed to detach from the tissue culture disc base. The changes in cell morphology were proportional to the increase of extract concentration (Figure 1).

Table 2. LAE and IAE content,

Test		Results	
Phytochemicals	Reagent	Observation	Conclusion
Flavonoid	wlister	Orange	Positive
Saponin	Forth	Clear and/or foamy	Positive
Tannin	FeCB	Yellow to orange	Positive
Alkaloid	Meyer	White ppt	Negative
Steroid	Liebermann-Burchard	Blackish blue	Positive
Terpenoid	-	Reddish brown	Positive




Figure 1. Effect of LAE and IAE treatment to cell morphology compared to control. Observation was done using inverted microscope, 100x magnification.

LAE and IAE inhibited NIH3T3 cells' growth, proportional to its concentration. IAE showed significantly higher inhibitory activity compared to EAL statistically ($p < 0.05$) (Figure 2). Cell viability obtained from the cytotoxicity testing could be calculated further to get IC_{50} value, showing its level of cytotoxicity. LAE and IAE exhibited low toxicity, giving the IC_{50} values in the range of 400-500 ppm (Table 3).

Cell Proliferation Assay

Citotoxicity assay showed that low concentration of LAE ($< 50 \mu\text{g/mL}$) could increase cell viability (Figure 3). Proliferation assay was conducted to confirm that the percentage of living cells was above 100%. LAE with the concentration of 10 and 25 ppm could increase fibroblast cells proliferation kinetics compared to control. In considerably high concentration (50 and 125 $\mu\text{g/}$




Figure 2. LAE and IAE inhibited NIH3T3 fibroblast cells growth. As much as 3×10^3 cells/well were treated with LAE and IAI, and cell viability were measured with MTT assay.

Table 3. Cytotoxicity potential of LAE and IAE on NIH3T3 cells.

Sample	IC ₅₀ (µg/mL)
LAE	529
IAE	463

mL), the increase of proliferation kinetics was also observed to occur up to the 48th hour, followed by decrease in the 72nd hours.

H₂O₂ Induction and Recovery

The purpose of stress induction using H₂O₂ was to observe the continuous protective effect of LAE and IAE on fibroblast cells against H₂O₂. Exposure was set to be 3 days to mimic chronic stress. LAE and IAE could increase fibroblast cells' viability pretreated with H₂O₂ stress induction. IAE treatment resulted in greater increase of cell viability compared to EAL in all tested concentrations (Figure 4). Thus, both extracts have potency as cell protector against H₂O₂ exposure.

LAE and IAE was proved to have cytoprotective activity on NIHT3T cells continuously against 100 µM H₂O₂ oxidative stress exposure given consecutively for 3 days, with the concentration of 50 and 125 µg/mL, giving

better protective activity compared to 352 µM vitamin C as positive control (*p*<0.05). To confirm the cytoprotective effect observed, recovery test on fibroblast cells against oxidative stress exposure needs to be conducted. It was aimed to reveal whether the extract could quickly protect NIHT3T cells undergo acute oxidative stress (1 day) treatment.

Recovery testing showed that % cell viability of cells treated with 50-500 ppm LAE gave greater effect compared to negative control, while IAE-treated cells showed higher % cell viability in all treatment concentration compared to negative controls (H₂O₂) (Figure 5). Vitamin C was used as positive control as its antioxidant activity has been proven. It has the ability to neutralize reactive oxygen, such as hydrogen peroxide. During the process, vitamin C itself turns into radical monohydroascorbate that will undergo reduction enzymatically by reduced glutation (GSH) catalyzed by glutation peroxidase back into vitamin C, yielding oxidized glutation (GSSG), and nonenzymatically via the reaction between two mono dehydrocorbate molecules yielding ascorbate and dehydroascorbate that are not radicals (Bender, 2009).




Figure 3. Fibroblast cells growth rate, time (hours vs absorbance). The test was conducted on fibroblast cells and observation was done after 0, 24, 48, and 72 hours.




Figure 4. Effect of LAE and IAE and on NIH3T3 fibroblast cells exposed with H₂O₂. Cells were treated with 50, 125, 250, 500, and 1000 ppm and 325 μM ascorbic acid for 1 hour prior to oxidative stress exposure using 100 μM H₂O₂ for 3 consecutive days. Statistical analysis showed significant difference ($p < 0.05$).

H₂O₂ gave oxidative stress to NIH3T3 fibroblast cells. It is non-radical oxygen derivative that will undergo a series of reactions yielding free radicals (Varh and Stroz, 2010). Free radicals are atoms or molecules that have one unpaired electron. As a result, free radicals possess high reactivity because of their tendency to pull electrons, hence they could convert another molecule into radicals as the result of losing or gaining an additional electron. Tissue damage caused by radical oxygen species (ROS) is called oxidative stress, while agents protecting the tissue from ROS are called antioxidants (Bender, 2009).

Apple contains substances that are expected to function as antioxidants. Phytochemicals expected to possess free-radical-stabilizing ability in LAE and IAE are flavonoids. Pertiwi, *et al.* (2016) revealed that apple peel contained flavanols, flavonoids, and quercetin. On the other hand, Nurcahyati (2015) showed that apple peel contains quercetin, a

dominant flavonol content in apple peel. Quercetin may act as an antioxidant and antiaging agent (Wasim and Farhan, 2010). Antioxidants could protect cells from damages caused by unstable molecules known as free radicals. Antioxidants are able to donate their electrons to free radicals. Thus from the study we may conclude that one of the secondary metabolites responsible for the cytoprotective ability is flavonoid, as one group of antioxidants. Not only flavonoids are known as antimutagenic and anticarcinogenic, they also act as antioxidants, anti-inflammatory agents, antihistamines, and inhibitors of LDL oxidation. LAE and IAE contain flavonoids that are antioxidants. Antioxidants have protective ability against oxidative stress and also inflammation. LAE and IAE with concentrations of 50 and 125 ppm showed better cytoprotective activity compared to vitamin C as a positive control ($p < 0.05$), that could possibly be because of their antioxidant content protecting cells from H₂O₂.




Figure 5. Recovery effect of LAE and IAE on NIH3T3 fibroblast cells exposed with H₂O₂. Cells were treated with 50, 125, 250, 500, and 1000 ppm LAE and IAE, and 352 μ M ascorbic acid for 1 hour prior to oxidative stress exposure using 100 μ M H₂O₂. Statistical analysis showed significant difference ($p < 0.05$).

exposure. The mechanism of flavonoids to protect the cells against free radicals was as follows: it acted as antioxidant scavenging water soluble free radicals to form a relatively stable radical and could stay for quite some time until it reacts with nonradical product (Bender, 2009). While with the concentration of 250, 500, and 1000 ppm, the decrease in % cell viability was observed, predicted to be the result of flavonoid content in LAE and LAI that underwent changes into oxidant radicals. Free radicals have two typical characteristics, that are high reactivity because of its tendency to attract electron and able to convert molecule into radical, that make it grouped into oxidants. In this case, substances in LAE and IAE could possibly undergo change into free radicals that cause the increase of total free radicals exposing the cells.

The substance that was predicted to cause the increase in cell viability was flavonol. Another substance that might increase cell viability was tannin, one of the polyphenols. Polyphenols in

apple could increase cell viability to up to 60% and in could efficiently protect yeast cells from H₂O₂ oxidative stress (Palermo, et al., 2012). Hence, the metabolites that could protect NIH3T3 fibroblast cells against chronic oxidative stress exposure was predicted to be flavonoids and polyphenols antioxidants.

CONCLUSION

LAE and IAE contained flavonoids, saponin, tannin, steroid, and terpenoid. LAE and LAI with the concentration of 50 and 125 ppm could protect NIH3T3 fibroblast cells against H₂O₂ oxidative stress exposure better than ascorbic acid.

REFERENCES

- Atmaja, N.S., Marwiyah and Setyowati, E., 2012, Pengaruh Kosmetika Anti Aging Wajah Terhadap Hasil Perawatan Kulit wajah, *J. Beaut. Beaut. Health Edu.*, 1(1), 1-5.

- Bender, D.A., 2009, Free Radicals and Antioxidant Nutrients. In: Murray K., Bender, D.A., Botham, K.M., Kennely, P.J., Rodwell, V.W., Weil, P.A., editors. *Harper's Illustrated Biochemistry*, 28th edition, New York: Mc Graw Hill Lange, p.482-486.
- Curtis, R., Geesaman, B.J. and DiStefano, P.S., 2005, Aging and Metabolism: Drug Discovery Opportunities, *Nat. Rev. Drug Discov.*, 4(7): 569-580.
- Gilchrist, B.A., Fitzpatrick, A.Z., Eisen, I.M., Freedberg and Austen, K.F., 2003, *Aging of Skin In Dermatology in General Medicine, Vol2*, 6th Edition. New York: Mc Graw-Hill.
- Juranovic, C.I., Zeiner, M., Kröppl, M. and Stingeder, G., 2011, Comparison of Sample Preparation Methods for the ICP-AES Determination of Major and Minor Elements in Clarified Apple Juices, *Microchem. J.*, 99(2): 364-369.
- Kujawa, D.B., Sylwia, C., Jan, O. and Halina, K., 2011, Extract From Apple Leaves and Fruits as Effective Antioxidants, *J. Med. Plant. Res.*, 5(11), 2339-2347.
- Getoff, N., 2007. Anti-aging and Aging Factors in Life. The Role of Free Radicals, *Radiat. Phys. Chem.*, 76(2007), 1577-1586.
- Nurchayati, E., 2014, *Khasiat dan Manfaat Dahsyatnya Kulit Apel Untuk Kesehatan Dan Penyembuhan*, Jakarta: Jendela Sehat.
- Palermo, V., Mattivi, F. and Mazzoni, C., 2012, Apple Can Act as Anti-aging on Yeast Cells, *Oxid. Med. Cell Longev.*, 2012(2012), 491759. doi: 10.1155/2012/491759.
- Pertiwi, R.D., Cut E.Y. dan Nanda F.P., 2016, Uji Aktivitas Antioksidan Ekstrak Etanol Limbah Kulit Apel (*Malus domestica* Borkh) Terhadap Radikal Bebas DPPH (2,2-Diphenyl-1-Picrylhydrazil), *Jurnal Ilmiah Manuntung*, 2(1), 81-92.
- Varh, L.G. and Stroz, P., 2010, Reactive Oxygen Species in Cancer, *Free Radic. Res.*, 44(5), 479-496.
- Wasim and Farhan, A., 2010, *Isolasi & Identifikasi Golongan Flavonoid Daun Dendang Gendis (Clinacanthus nutans)*, Essay, Universitas Negeri Yogyakarta, Yogyakarta
- Yuniarto P.F., Rahayu, E.S.R. and Ekowati, D., 2012, Optimasi Formula Gel Buah Apel Hijau (*Pyrus malus* L.) sebagai Antioksidan dengan Kombinasi Basis Carbopol 940 dan Gliserin secara Simplex Lattice Design, *Jurnal Farmasi Indonesia*, 11(2), 130-138.